


RESEARCH ARTICLE

Open Access

Arabidopsis paralogous genes *RPL23aA* and *RPL23aB* encode functionally equivalent proteins



Wei Xiong^{1,2}, Xiangze Chen¹, Chengxin Zhu¹, Jiancong Zhang¹, Ting Lan^{1,2}, Lin Liu¹, Beixin Mo^{1*}  and Xuemei Chen^{3*}

Abstract

Background: In plants, each ribosomal protein (RP) is encoded by a small gene family but it is largely unknown whether the family members are functionally diversified. There are two *RPL23a* paralogous genes (*RPL23aA* and *RPL23aB*) encoding cytoplasmic ribosomal proteins in *Arabidopsis thaliana*. Knock-down of *RPL23aA* using RNAi impeded growth and led to morphological abnormalities, whereas knock-out of *RPL23aB* had no observable phenotype, thus these two *RPL23a* paralogous proteins have been used as examples of ribosomal protein paralogues with functional divergence in many published papers.

Results: In this study, we characterized T-DNA insertion mutants of *RPL23aA* and *RPL23aB*. A rare non-allelic non-complementation phenomenon was found in the F1 progeny of the *rpl23aa* X *rpl23ab* cross, which revealed a dosage effect of these two genes. Both *RPL23aA* and *RPL23aB* were found to be expressed almost in all examined tissues as revealed by GUS reporter analysis. Expression of *RPL23aB* driven by the *RPL23aA* promoter can rescue the phenotype of *rpl23aa*, indicating these two proteins are actually equivalent in function. Interestingly, based on the publicly available RNA-seq data, we found that these two *RPL23a* paralogues were expressed in a concerted manner and the expression level of *RPL23aA* was much higher than that of *RPL23aB* at different developmental stages and in different tissues.

Conclusions: Our findings suggest that the two *RPL23a* paralogous proteins are functionally equivalent but the two genes are not. *RPL23aA* plays a predominant role due to its higher expression levels. *RPL23aB* plays a lesser role due to its lower expression. The presence of paralogous genes for the *RPL23a* protein in plants might be necessary to maintain its adequate dosage.

Keywords: Ribosome, Ribosomal protein, *RPL23a*, Gene dosage, Parologue, Functional specialization

* Correspondence: bmo@szu.edu.cn; xuemei.chen@ucr.edu

¹Guangdong Provincial Key Laboratory for Plant Epigenetics, Longhua Bioindustry and Innovation Research Institute, College of Life Sciences and Oceanography, Shenzhen University, Shenzhen 518060, Guangdong, China

³Department of Botany and Plant Sciences, Institute of Integrative Genome Biology, University of California, Riverside, CA 92521, USA

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Ribosomes are responsible for protein synthesis in all living cells. A single ribosome is a ribonucleoprotein complex formed from a large and a small subunit. In plants, the large ribosomal subunit is composed of 28S, 5.8S and 5S rRNAs together with 48 RPL (Ribosomal Protein of Large subunit) proteins, whereas the small subunit is composed of 18S rRNA and 33 RPS (Ribosomal Protein of Small subunit) proteins [1, 2]. In *E. coli*, genes encoding RPs are arranged in about 20 operons, with approximately half of the genes mapping to a single locus [3, 4]. In mammals, although there are about 2000 sequences which may encode RPs, most of them are predicted to be pseudogenes, and most functional RPs are encoded by a single copy [5]. In yeast *Saccharomyces cerevisiae*, about 75% of the RPs are encoded by gene families with more than one member [6]. Although substantially functional redundancy was found between paralogous RP genes in yeast, some paralogous RP genes were reported to have non-redundant functions [6–8].

Plants have even more gene members encoding a single RP than yeast [9]. In *Arabidopsis thaliana*, RP paralogues share 65 to 100% amino acid sequence identity [9]. Assessment of cognate EST (expressed sequence tag) numbers of RP genes suggested that RP gene family members were differentially expressed in *Arabidopsis* [9]. Microarray data also revealed that transcripts of RP genes within the same family were accumulated at different levels in *Arabidopsis* [10]. Under various stimuli, while the transcript levels for most RP genes remain unchanged, some RP genes show altered expression levels [10]. Many studies have investigated the phenotypic consequence of absent or reduced expression of a single RP paralogue in *Arabidopsis*. Disruptions in any one of the RP protein genes, *RPL3A*, *RPL8A*, *RPL19A*, *RPL23C*, *RPL40B*, and *RPS11A* is embryo lethal [11]. Less severe phenotypes were reported for mutations in several other RPs [11]. Morphological changes of the first two true leaves from the spatulate wild type shape to a pointed shape were found in mutants of some RP genes, including *RPL5A*, *RPL5B*, *RPL9C*, *RPL10aB*, *RPL24B*, *RPL28A*, *RPS13B*, and *RPS18A* [12–16]. Despite these studies on RPs, it remains unknown why RPs are encoded by paralogues in plants or whether RP paralogues have specialized functions.

In *Arabidopsis*, the *RPL23a* family consists of two members (*RPL23aA* and *RPL23aB*) that encode proteins with 95% amino acid identity. Both *RPL23aA* and *RPL23aB* genes are transcribed and translated, and protein products of either paralogue can be incorporated into the cytoplasmic ribosome [17, 18]. Knock-down of the *RPL23aA* gene through RNAi results in severe developmental defects, whereas knock-down, or even knock-out, of *RPL23aB* has no obviously phenotypic

consequences [19], which could be the basis for the argument that *RPL23aA* and *RPL23aB* had specialized functions [11, 19–21].

With the general question of why plant RPs are encoded by paralogous genes in mind, we sought to study the functional relationship between *RPL23aA* and *RPL23aB*. With T-DNA insertion mutants in *RPL23aA* and *RPL23aB*, we found a rare non-allelic non-complementation phenomenon, indicating that these two *RPL23a* genes are dosage dependent genes. We showed that expression of *RPL23aB* driven by the *RPL23aA* promoter can rescue the phenotype of *rpl23aa*, demonstrating that RPL23aA and RPL23aB proteins are functionally equivalent. Furthermore, interrogation of RNA-seq data from several developmental stages and in different organs showed that although the level of *RPL23aA* transcripts was much higher than that of *RPL23aB*, the fluctuations in expression of the two genes were well matched, suggesting that these two genes were coordinately regulated. These results revealed that duplicated *RPL23a* genes contribute to ribosome dosage necessary for plant growth and development. Our results do not contradict prior studies showing that *RPL23aA* plays a dominant role in plant growth and development, but reveal that the *RPL23aA* dominance resides in its higher expression level rather than functional specialization of the protein.

Methods

Plant material and growth conditions

Arabidopsis thaliana wild type Columbia-0 (Col-0) and the T-DNA insertion lines, *SALK_005448* (named here *rpl23aa*) and *SAIL_597_B08* (named here as *rpl23ab*), were obtained from the *Arabidopsis* Biological Resource Center (ABRC). Seeds were first treated for 2 min in 75% ethanol, then treated for 6 min in commercial bleach and rinsed at least 3 times with sterile distilled water. Solid medium consisted of 2.2 g/L Murashige and Skoog basal salt mixture (Phyto Tech Labs), 10 g/L sucrose, and 8 g/L agar. pH was adjusted to 5.6 with KOH before autoclaving. When required, BASTA (GOLDBIO) was added at a final concentration of 125 µg/L. Seeds were sown in a water suspension, using a 1.5 mL pipette tip, in 150 mm Petri dishes filled with 120 ml of solid culture medium, at a density of 150 regularly spaced seeds per plate. Once inoculated, the Petri dishes were sealed with Micropore Scotch 3M surgical tape, which prevented contamination but allowed gaseous exchange, and placed in 4 °C for 24 h. Growth was allowed to proceed at 22 °C in Percival tissue culture chambers under long day conditions (16 h light and 8 h dark). 10-day seedlings were then transplanted to pots containing a 1:2:2 mixture of perlite, vermiculite and soil at 22 °C under long day conditions from a combination of

incandescent and fluorescent lamps (10,000 lx). Plants were watered twice a week with nutrient solution.

RNA isolation and RT-PCR

50 mg seedlings from 14-day-old Col-0, *rpl23aa*, and *rpl23ab* were harvested and immediately frozen in liquid nitrogen. RNA was extracted using RNAiso Plus (TAKARA BIO INC). In the elution step, RNA was resuspended in DEPC-treated water. cDNA was obtained by reverse transcription of 1 µg of RNA with the PrimeScript™ RT reagent Kit with gDNA Eraser (TAKARA BIO INC).

Plasmid construction and generation of transgenic plants

In order to construct the pRPL23aA::RPL23aA and pRPL23aB::RPL23aB plasmids, a 3001 bp DNA fragment (including the promoter region) of *RPL23aA* (AT2G39460) and a 2016 bp DNA fragment (including the promoter region) of *RPL23aB* (AT3G55280) were amplified from Col-0 genomic DNA using Phusion polymerase (Thermo Scientific). The primers used are shown in Table S1 (Additional file 12). The amplified DNA sequences were cloned in pEG301 [22] to result in pRPL23aA::RPL23aA and pRPL23aB::RPL23aB. The plasmids were used to transform *rpl23aa*. For pRPL23aA::RPL23aB construction, the promoter region (about 1.5 kb) of *RPL23aA* plus the coding region of *RPL23aB* were synthesized by a commercial company (GENEWIZ SuZhou), then the synthesized DNA fragment was sequenced and was cloned in pEG301. The promoter regions of *RPL23aA* (about 1.5 kb) and *RPL23aB* (about 1.5 kb) were cloned into pMDC162 [22] to generate the plasmids pRPL23aA::GUS and pRPL23aB::GUS, which were then used to transform Col-0 plants. Floral dip transformation was performed as described by Clough and Bent [23]. T1 transgenic plants were screened on solid 1/2 Murashige & Skoog (MS) medium with 25 mg/L Hygromycin B or 0.002% BASTA and verified by PCR. GUS staining was carried out with plants in the T2 generation.

GUS staining assay

8-days-old seedlings and 36-days-old inflorescences, immature and mature flowers, immature and mature siliques of Col-0, pRPL23aA::GUS and pRPL23aB::GUS were subjected to histochemical GUS staining according to the standard protocol [24].

Transcripts profiling

RNA-seq data was obtained from a public website (<http://travadb.org/browse/DeSeq/>), and the average value of normalized absolute read counts from two biological replicates was extracted. We also downloaded the original RNA-seq data of *A. thaliana* different organs and developmental stages from NCBI Sequence Read

Archive (project ID PRJNA314076 for samples except meristem and project ID PRJNA268115 for the meristem samples). The RPKM (Reads Per Kilobase per Million mapped reads) value of *RPL23aA* (AT2G39460), *RPL23aB* (AT3G55280), and *ACT2* (AT3G18780) were calculated. Our calculated RPKM value is consistent with the value of normalized absolute read counts obtained from the public website (<http://travadb.org/browse/DeSeq/>).


Polysome profiling

Polysome profiling was performed as described by Mustroph et al. [25]. Briefly, 2 g of 14-day-old seedlings were collected and ground to a fine powder using sufficient liquid nitrogen, and the powder was resuspended in 8 mL of ice-cold polysome extraction buffer by gentle shaking. The lysate was incubated on ice for 10 min and centrifuged at 4 °C, 16,000 × g for 15 min. The supernatant was filtered through Miracloth and centrifuged at 4 °C, 16,000 × g for another 15 min. The supernatant was gently transferred to the top of a sucrose cushion and then centrifuged at 4 °C, 50,000 r.p.m. for 3 h to obtain the polysome pellet. The pellet was resuspended in ice-cold resuspension buffer and loaded onto a 4.5 mL sucrose gradient (20–60% w/v) for fractionation of polysomes by ultracentrifugation, after which the sucrose gradient was pumped through a UV detector and absorbance at 254 nm was recorded.

Results

Characterization of *rpl23aa* and *rpl23ab* mutants

The Arabidopsis genome contains two *RPL23a* paralogous genes *RPL23aA* (At2g39460) and *RPL23aB* (At3g55280), which encode proteins with 95% amino acids identity (see Additional file 1). We acquired T-DNA insertion lines of *RPL23aA* and *RPL23aB*, namely *SALK_005448* and *SAIL-597-B08*, respectively (hereafter referred to as *rpl23aa* and *rpl23ab*). PCR-genotyping confirmed that both *rpl23aa* and *rpl23ab* are homozygous T-DNA insertion alleles (see Additional file 2). Sequencing results revealed that *rpl23aa* contains a T-DNA insertion in the 3' UTR region, 10 bp downstream of the stop codon of the *RPL23aA* gene (Fig. 1a), while *rpl23ab* contains a T-DNA insertion in the second exon of *RPL23aB* (Fig. 1b). A semi-quantitative RT-PCR assay was used to detect transcripts from *RPL23aA* and *RPL23aB* in these T-DNA lines. As shown in Fig. 1c, the 3' region around the stop codon of the *RPL23aA* mRNA was disrupted in the mutant. Because the majority of the *RPL23aA* mRNA from the T-DNA line was intact, we suspect that *SALK_005448* is a hypomorphic allele. *rpl23ab* is likely a null mutant, because no *RPL23aB* mRNA was detected (Fig. 1d). Absence of dosage compensation by *RPL23aA* in Arabidopsis was reported



following loss of *RPL23aB* [26]. As shown in Fig. 1d, there is also no dosage compensation by *RPL23aB* in the *rpl23aa* mutant.

The *rpl23aa* mutant exhibits pleiotropic defects, including pointed leaves, retarded root growth, and reduced plant size (Fig. 2b). These phenotypes are similar

to those of a previously reported RNAi line [19]. An incompletely penetrant tricotyledon phenotype (less than 5% of the total population) was observed in *rpl23aa* mutant plants (see Additional file 3). However, we didn't observe appreciable defects in terms of growth rate, morphology, or flowering in the *rpl23ab* mutant (Fig.




Fig. 2 Plant phenotypes. 14-day-old plants of (a) Col-0, (b) *rpl23aa*, (c) *pRPL23aA::RPL23aA-HA/rpl23aa*, (d) *rpl23ab*. *rpl23aa* exhibits pleiotropic defects, including pointed leaves, retarded root growth, and reduced plant size; *pRPL23aA::RPL23aA-HA* fully rescued the morphological defects of *rpl23aa*; *rpl23ab* had no observable phenotype. Size bar, 2 mm

2d), which is consistent with published work [26]. We amplified genomic DNA encompassing the promoter plus the coding region of *RPL23aA* from wild-type plants and fused it to the sequence encoding the HA epitope tag. When this transgene was introduced into *rpl23aa*, the developmental defects were fully rescued (Fig. 2c), suggesting that dysfunction of *RPL23aA* was responsible for the developmental defects in *rpl23aa*.

***RPL23aA* and *RPL23aB* are dosage-dependent genes**

In order to study the genetic interaction between *RPL23aA* and *RPL23aB*, we crossed *rpl23aa* with *rpl23ab*. To our surprise, the doubly heterozygous plants (*RPL23aA/rpl23aa; RPL23aB/rpl23ab*) in the F1 progeny all have pointed first true leaves (Fig. 3b). Siliques of the doubly heterozygous plants are much shorter than siliques of *rpl23aa* or *rpl23ab* (Fig. 3i). We dissected siliques from *RPL23aA/rpl23aa; RPL23aB/rpl23ab* plants and found many aborted ovules (Fig. 3g and h). An F2 population was generated by selfing the above F1 plants. We genotyped 144 F2 plants but did not find double homozygous (*rpl23aa/rpl23aa; rpl23ab/rpl23ab*)

plants. In fact, we did not even detect any genotypes with a single functional allele from either gene (*RPL23aA/rpl23aa; rpl23ab/rpl23ab* or *rpl23aa/rpl23aa; RPL23aB/rpl23ab*) (Table 1), although these genotypes are collectively expected to appear in 31.25% (5 out of 16) of the F2 plants. We suspected that this non-allelic non-complementation phenomenon between *rpl23aa* and *rpl23ab* is probably due to gene dosage effects.

***RPL23aA* and *RPL23aB* genes have similar expression patterns**

In order to investigate the expression patterns of *RPL23aA* and *RPL23aB* genes, we fused the promoter regions of *RPL23aA* and *RPL23aB* genes to the GUS reporter and generated transgenic plants in the Col-0 background. GUS staining of 14 *pRPL23aA::GUS* and 5 *pRPL23aB::GUS* independent transgenic lines uncovered a ubiquitous expression pattern for both genes with particularly intense GUS staining in young and actively proliferating tissues, such as developing leaves, floral buds and root apices (Fig. 4). Similar expression patterns of




Fig. 3 The non-allelic non-complementation phenomenon between *rpl23aa* and *rpl23ab*. 9-day-old plants of (a) *rpl23aa* × Col-0 (F1 generation), (b) *rpl23aa* × *rpl23ab* (F1 generation). Dissected mature siliques from (c) Col-0, (d) *rpl23aa*, (e) *rpl23ab*, (f) *rpl23aa* × Col (F1 generation), (g) *rpl23aa* × *rpl23ab* (F1 generation), (h) *rpl23ab* × *rpl23aa* (F1 generation). (i) The length of mature silique from *rpl23ab*, *rpl23aa*, and the double heterozygote (double het). Arrowheads indicate aborted embryos. Size bar, 5 mm

RPL23aA and *RPL23aB* support our hypothesis that the non-allelic non-complementation phenomenon between these two genes is the consequence of overlap in expression (and function) of *RPL23aA* and *RPL23aB* in the same cells.

RPL23aA and RPL23aB proteins are functionally equivalent

It has been reported that some paralogous ribosomal proteins have evolved specialized functions in yeast [6]. As mentioned above, dysfunction of *RPL23aA* results in

severe developmental defects, whereas knock-out of *RPL23aB* has no phenotypic consequences in Arabidopsis ([19, 26] and this study). It's natural to assume that these two paralogous ribosomal proteins have undergone functional specialization.

We designed gene complementation experiments to explore whether *RPL23aA* and *RPL23aB* have distinct functions. If *RPL23aA* and *RPL23aB* have specialized functions, *RPL23aB* is not expected to complement the *rpl23aa* mutation. We fused the promoter regions of *RPL23aA* to the coding region of *RPL23aB*. The

Table 1 Genetic interactions between *rpl23aa* and *rpl23ab*

Genotype	First leaf phenotype	
	Pointed	Normal
<i>RPL23aA/RPL23aA RPL23aB/RPL23aB</i>	0	15
<i>RPL23aA/RPL23aA RPL23aB/rpl23ab</i>	0	37
<i>RPL23aA/RPL23aA rpl23ab/rpl23ab</i>	0	19
<i>RPL23aA/rpl23aa RPL23aB/RPL23aB</i>	0	26
<i>RPL23aA/rpl23aa RPL23aB/rpl23ab</i>	38	0
<i>RPL23aA/rpl23aa rpl23ab/rpl23ab</i>	0	0
<i>rpl23aa/rpl23aa RPL23aB/RPL23aB</i>	9	0
<i>rpl23aa/rpl23aa RPL23aB/rpl23ab</i>	0	0
<i>rpl23aa/rpl23aa rpl23ab/rpl23ab</i>	0	0

rpl23aa and *rpl23ab* were crossed and the F₂ plants were subjected to genotyping at the *RPL23aA* and *RPL23aB* loci. Leaf phenotype of the plants was classified into pointed or normal. Primers for genotyping are listed in Table S1 (Additional file 12)

pRPL23aA:RPL23aB transgene was introduced into *rpl23aa* plants, and 21 independent *pRPL23aA:RPL23aB* transgene lines were obtained, among which 15 lines rescued the phenotype of *rpl23aa* (Fig. 5 and Additional file 4), indicating that *RPL23aA* and *RPL23aB* have equivalent function. The *pRPL23aB:RPL23aB* transgene was also introduced into *rpl23aa* plants, and 8 out of 15 independent, homozygous transgenic lines exhibited near wild type morphology (Fig. 5 and Additional file 4). However, a portion (about 2%) of the transgenic plants of each line exhibited the tricotyledon phenotype (see Additional file 5). Thus, the *pRPL23aB:RPL23aB*

transgene can largely but not fully rescue the phenotype of *rpl23aa*.

RPL23aA and *RPL23aB* genes are transcribed in a concerted manner with higher expression levels of *RPL23aA* than *RPL23aB*

Since the above results indicated that *RPL23aA* and *RPL23aB* proteins have equivalent function, we suspected that the difference in phenotype between *rpl23aa* and *rpl23ab* is due to the difference in the expression levels of these two genes. The expression of *RPL23aA* may be much higher than *RPL23aB*, so the impacts on ribosomes by the *rpl23aa* mutation are higher than the *rpl23ab* mutation thus leading to much severe morphological defects. We compared the transcript levels of *RPL23aA* and *RPL23aB* at different developmental stages and in different organs by analyzing published RNA-seq data [27]. As shown in Fig. 6 and Figure S6 (Additional file 6), transcript levels of *RPL23aA* are much higher than those of *RPL23aB* at all developmental stages and in all the examined tissues. Strikingly, the spatial and temporal patterns of expression of these two paralogous genes are well matched, suggesting that they are similarly regulated at differently developmental stages in all examined tissues. *ACT2*, which is a house keeping gene, was included as a control. Transcript levels of *RPL23aB* are higher than *ACT2* in some organs, and total amount of *RPL23a* transcripts is much higher than *ACT2* in most examined organs (Fig. 6c, e), indicating that RPs are in great demand for plant development.




Fig. 4 Promoter-GUS reporter analysis of *RPL23aA* and *RPL23aB*. **a, b, c** Seedling. **d, e, f** Inflorescences. **g, h, i** Immature and **j, k, l** mature flowers. **m, n, o** Immature and **p, q, r** mature siliques. Pictures were taken at 8 days **a-c** and 36 days **d-r**. Size bar, 5 mm




The polysome/monosome ratio is elevated in the *rpl23a* mutants

In order to evaluate the effects of the *rpl23aa* and *rpl23ab* mutations on polysomes, we fractionated total ribosomes by ultracentrifugation through sucrose density gradients. The polysome profiles from plants of various genotypes are shown in Figure S7 (Additional file 7). To our surprise, the ratio between polysomes and the monosome is obviously increased in *rpl23aa* and slightly increased in *rpl23ab* compared to wildtype. The increase in the polysome/monosome ratio in *rpl23aa* was largely rescued by both *pRPL23aA::RPL23aA* and *pRPL23aA::RPL23aB* transgenes, whereas the polysome/monosome ratio in *pRPL23aB::RPL23aB/rpl23aa* transgenic plants is higher than wildtype and lower than *rpl23aa*. The changes in ribosomal profile of *rpl23aa* and *rpl23ab* indicate that the overall translation state is altered. The higher polysome levels could reflect higher rates of translation or defects in translation, such as slower elongation. While the molecular basis of the higher polysome levels is unknown, the stronger effect of the *rpl23aa* mutation is consistent with the dominant role of *RPL23aA* over *RPL23aB* as suggested by expression levels and mutant phenotypes.

Discussion

Some of the paralogous RPs are identical in amino acid sequences such as *RPL36aA* and *RPL36aB*, but many of the paralogues display sequence variations and are differentially expressed during development. The presence of multiple gene members for each RP in plants might be necessary to maintain adequate RP doses or to maintain some degree of ribosome heterogeneity and functional specialization.

In this study, we characterized the *RPL23a* gene family containing two highly homologous family members. The hypomorphic T-DNA insertion allele of *RPL23aA* exhibits pleiotropic defects. However, knock-out of *RPL23aB* has no appreciable phenotypic impacts. We crossed mutants of *RPL23aA* and *RPL23aB* and found a non-allelic non-complementation phenomenon in their F1 progeny. This phenomenon is also found in other RP coding gene families such as *RPL5* [28], *RPL36a* [29], and *RPS6* [30]. However, mutations in the paralogues within *RPL5*, *RPL36a*, and *RPS6* families caused almost the same phenotype, indicating that the paralogues are functionally equivalent. In the case of the *RPL23a* family, phenotypes of the single mutants suggest unequal functions of the two paralogues. The non-allelic non-complementation phenomenon may be due to a dosage



problem - reduced dosage at one of the paralogues still supports the wild phenotype but simultaneous reduction of dosage at both paralogues could not sustain the wild phenotype. For the dosage effect hypothesis to be true, there must be at least some overlap in the expression of the gene family members. Indeed, promoter-GUS experiments demonstrated that both *RPL23aA* and *RPL23aB* were ubiquitously expressed.

Phenotypical differences between members of an RP within a family might result from diversification of protein function or variation in levels and patterns of expression. We demonstrated that *RPL23aA* and *RPL23aB* proteins had equal function, as expression of *RPL23aB* driven by the *RPL23aA* promoter could rescue the phenotype of the *rpl23aa* mutant. We found that the expression level of *RPL23aA* was much higher than that of *RPL23aB* according to the publicly available RNA-seq data. Thus, the difference in expression levels might be the reason why disruption of *RPL23aA* and *RPL23aB*

had different consequences. It is interesting that despite the difference in expression levels, the temporal and spatial patterns of expression of the two paralogous genes were nearly identical. These results suggested that *RPL23aA* and *RPL23aB* genes are transcribed in a coordinated manner. Posttranscriptional and translational regulation may also play a role in *RPL23aA* and *RPL23aB* expression [31]. Subcellular localization specialization could be another factor that causes differences in functional effects between paralogous RPs [32]. Previous studies revealed that both of *RPL23aA* and *RPL23aB* are targeted to the nucleolus with *RPL23aA* targeting being a bit more efficient than *RPL23aB* [10, 19]. Targeting of RP to the nucleolus is an essential step in eukaryotic ribosome biogenesis [33, 34], so the efficiency of *RPL23aA* assembly into ribosomes may be higher than that of *RPL23aB*. Although posttranscriptional differences between *RPL23aA* and *RPL23aB* may exist, the fact that expression of *RPL23aB* with the

RPL23aA promoter rescues the *rpl23aa* phenotypes indicates that differences in expression level underlie the different functional contributions of the paralogues as exemplified by the single mutant phenotypes.

There are at least four possible consequences of a RP disruption: (1) ribosome insufficiency, (2) non-functional ribosomes, (3) partially dysfunctional ribosomes, and (4) loss of the extraribosomal function of the RP [35]. Polysome profiling results revealed that the polysome/monosome ratio is elevated in the *rpl23a* mutants, which suggested global translational alteration. The exact nature of the alteration remains unknown and will be investigated in the future.

Conclusions

Ribosomal protein RPL23a paralogues (*RPL23aA* and *RPL23aB*) have been used as examples of paralogues with functional divergence in many published papers. In this study, our findings provided four convincing evidences demonstrating duplicated *RPL23a* genes actually have redundant function (without functional specialization), thus are necessary to provide a threshold dose: 1) The non-allelic non-complementation phenomenon between *rpl23aa* and *rpl23ab* suggests *RPL23aA* and *RPL23aB* are dosage dependent genes; 2) *RPL23aA* and *RPL23aB* genes are expressed in the same tissues; 3) *RPL23aB* could rescue the phenotype of *rpl23aa*, demonstrating *RPL23aA* and *RPL23aB* protein have equal function; 4) *RPL23aA* and *RPL23aB* genes are transcribed in a concerted manner with higher expression levels of *RPL23aA* than *RPL23aB*. Our findings suggest that the two paralogous *RPL23a* proteins have equivalent function and the presence of multiple genes for individual RPs in plants might be necessary to maintain adequate ribosome dosage at least for some ribosomal protein families.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s12870-020-02672-1>.

Additional file 1: Figure S1. Amino acid sequence alignment between *RPL23aA* and *RPL23aB*.

Additional file 2: Figure S2. Genotyping of *rpl23aa* and *rpl23ab*.

Additional file 3: Figure S3. Images of wild type and *rpl23aa* plants.

Additional file 4: Figure S4. Lengths of mature siliques and numbers of ovules in mature siliques.

Additional file 5: Figure S5. Images of *pRPL23aB:RPL23aB rpl23aa* plants.

Additional file 6: Figure S6. Transcript profiles of *RPL23aA* and *RPL23aB* in different organs.

Additional file 7: Figure S7. Polysome profiles of Col-0 (black), *rpl23aa* (red), *rpl23ab* (green), *pRPL23aA:RPL23aA/rpl23aa* (yellow), *pRPL23aA:RPL23aB/rpl23aa* (blue), and *pRPL23aB:RPL23aB/rpl23aa* (purple).

Additional file 8: Figure S8. Full-length gel of Fig. 1c.

Additional file 9: Figure S9. Full-length gel of Fig. 1d.

Additional file 10: Figure S10. Full-length gel of Figure S2C.

Additional file 11: Figure S11. Full-length gel of Figure S2D.

Additional file 12: Table S1. Primers used in this work.

Abbreviations

RP: ribosomal protein; RPL: ribosomal protein of large subunit; RPS: ribosomal protein of small subunit; RPKM: Reads Per Kilobase per Million mapped reads

Acknowledgements

We thank Doctor Wenwen Kong and Doctor Yang Liu from Shenzhen University for their help with RNA-seq data analysis.

Authors' contributions

XC and BM designed experiments; WX, XZC, and CZ carried out experiments; WX, JZ, and TL analyzed the RNA-seq data; XW and LL analyzed experimental results; WX, XZC, XC and BM wrote the manuscript. All authors agree to be accountable for the content of the work.

Funding

The authors thank the funding from Guangdong Innovation Team Project (2014ZT055078), National Natural Science Foundation of China (31870287) and National Key R&D Program of China Grant (2019YFA0903902). The funding body played no role in the design of the study, the collection, analysis and interpretation of the data and in writing the manuscript.

Availability of data and materials

The original RNA-seq data of *A. thaliana* different organs and developmental stages were downloaded from NCBI Sequence Read Archive (project ID PRJNA314076 for samples except meristem and project ID PRJNA268115 for the meristem samples).

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Guangdong Provincial Key Laboratory for Plant Epigenetics, Longhua Bioindustry and Innovation Research Institute, College of Life Sciences and Oceanography, Shenzhen University, Shenzhen 518060, Guangdong, China. ²Key Laboratory of Optoelectronic Devices and Systems of Ministry of Education and Guangdong Province, College of Optoelectronic Engineering, Shenzhen University, Shenzhen, Guangdong 518060, China. ³Department of Botany and Plant Sciences, Institute of Integrative Genome Biology, University of California, Riverside, CA 92521, USA.

Received: 27 February 2020 Accepted: 23 September 2020

Published online: 08 October 2020

References

- Chang IF, Szick-Miranda K, Pan SQ, Bailey-Serres J. Proteomic characterization of evolutionarily conserved and variable proteins of Arabidopsis cytosolic ribosomes. *Plant Physiol.* 2005;137(3):848–62.
- Carroll AJ. The Arabidopsis cytosolic ribosomal proteome: from form to function. *Front Plant Sci.* 2013;4:32.
- Mager WH. Control of ribosomal-protein gene-expression. *Biochim Biophys Acta.* 1988;949(1):1–15.
- Wikstrom PM, Bjork GR. A regulatory element within a gene of a ribosomal-protein operon of Escherichia-Coli negatively controls expression by decreasing the translational efficiency. *Mol Gen Genet.* 1989;219(3):381–9.
- Balasubramanian S, Zheng DY, Liu YJ, Fang G, Frankish A, Carriero N, Robilotto R, Cayting P, Gerstein M. Comparative analysis of processed ribosomal protein pseudogenes in four mammalian genomes. *Genome Biol.* 2009;10(1):R2.
- Komili S, Farny NG, Roth FP, Silver PA. Functional specificity among ribosomal proteins regulates gene expression. *Cell.* 2007;131(3):557–71.

7. Deutschbauer AM. Mechanisms of haploinsufficiency revealed by genome-wide profiling in yeast. *Genetics*. 2005;169(4):1915–25.
8. Dean EJ, Davis JC, Davis RW, Petrov DA. Pervasive and persistent redundancy among duplicated genes in yeast. *PLoS Genet*. 2008;4(7):e1000113.
9. Barakat A, Szickmiranda K, Chang I, Guyot R, Blanc G, Cooke R, Delseny M, Baileyserres J. The Organization of Cytoplasmic Ribosomal Protein Genes in the Arabidopsis genome. *Plant Physiol*. 2001;127(2):398–415.
10. Savada RP, Bonham-Smith PC. Differential transcript accumulation and subcellular localization of Arabidopsis ribosomal proteins. *Plant Sci*. 2014;223:134–45.
11. Byrne ME. A role for the ribosome in development. *Trends Plant Sci*. 2009;14(9):512–9.
12. Pinon V, Etchells JP, Rossignol P, Collier SA, Arroyo JM, Martienssen RA, Byrne ME. Three PIGGYBACK genes that specifically influence leaf patterning encode ribosomal proteins. *Development*. 2008;135(7):1315–24.
13. Ito T, Kim GT, Shinozaki K. Disruption of an Arabidopsis cytoplasmic ribosomal protein S13-homologous gene by transposon-mediated mutagenesis causes aberrant growth and development. *Plant J*. 2000;22(3):257–64.
14. Yao Y, Ling QH, Wang H, Huang H. Ribosomal proteins promote leaf adaxial identity. *Development*. 2008;135(7):1325–34.
15. Nishimura T, Wada T, Yamamoto KT, Okada K. The Arabidopsis STV1 protein, responsible for translation reinitiation, is required for auxin-mediated gynoecium patterning. *Plant Cell*. 2005;17(11):2940–53.
16. Vanlijsebettens M, Vanderhaeghen R, Deblock M, Bauw G, Villarroel R, Vanmontagu M. An S18 ribosomal-protein gene copy at the Arabidopsis Pfl locus affects plant development by its specific expression in meristems. *EMBO J*. 1994;13(14):3378–88.
17. McIntosh KB, Bonham-Smith PC. The two ribosomal protein L23A genes are differentially transcribed in Arabidopsis thaliana. *Genome*. 2005;48(3):443–54.
18. Carroll AJ, Heazlewood JL, Ito J, Millar AH. Analysis of the Arabidopsis cytosolic ribosome proteome provides detailed insights into its components and their post-translational modification. *Mol Cell Proteomics*. 2008;7(2):347–69.
19. Degenhardt RF, Bonham-Smith PC. Arabidopsis ribosomal proteins RPL23aA and RPL23aB are differentially targeted to the nucleolus and are disparately required for Normal development. *Plant Physiol*. 2008;147(1):128–42.
20. Xue SF, Barna M. Specialized ribosomes: a new frontier in gene regulation and organismal biology. *Nat Rev Mol Cell Bio*. 2012;13(6):355–69.
21. Whittle CA, Krochko JE. Transcript profiling provides evidence of functional divergence and expression networks among ribosomal protein gene paralogs in Brassica napus. *Plant Cell*. 2009;21(8):2203–19.
22. Earley KW, Haag JR, Pontes O, Opper K, Juehne T, Song K, Pikaard CS. Gateway-compatible vectors for plant functional genomics and proteomics. *Plant J*. 2006;45(4):616–29.
23. Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J*. 1998;16(6):735–43.
24. Jefferson RA, Kavanagh TA, Bevan MW. Gus fusions - Beta-Glucuronidase as a sensitive and versatile gene fusion marker in higher-plants. *EMBO J*. 1987;6(13):3901–7.
25. Mustroph A, Juntawong P, Bailey-Serres J. Isolation of plant polysomal mRNA by differential centrifugation and ribosome immunopurification methods. *Methods Mol Biol*. 2009;553:109–26.
26. Degenhardt RF, Bonham-Smith PC. Transcript profiling demonstrates absence of dosage compensation in Arabidopsis following loss of a single RPL23a paralog. *Planta*. 2008;228(4):627–40.
27. Klepikova AV, Kasianov AS, Gerasimov ES, Logacheva MD, Penin AA. A high resolution map of the Arabidopsis thaliana developmental Transcriptome based on RNA:seq profiling. *Plant J*. 2016;88(6):1058.
28. Fujikura U, Horiguchi G, Ponce MR, Micol JL, Tsukaya H. Coordination of cell proliferation and cell expansion mediated by ribosome-related processes in the leaves of Arabidopsis thaliana. *Plant J*. 2009;59(3):499–508.
29. Casanova-Saez R, Candela H, Micol JL. Combined haploinsufficiency and purifying selection drive retention of RPL36a paralogs in Arabidopsis. *Sci Rep*. 2014;4:4122.
30. Creff A, Sormani R, Desnos T. The two Arabidopsis RPS6 genes, encoding for cytoplasmic ribosomal proteins S6, are functionally equivalent. *Plant Mol Biol*. 2010;73(4–5):533–46.
31. McIntosh KB, Degenhardt RF, Bonham-Smith PC. Sequence context for transcription and translation of the Arabidopsis RPL23aA and RPL23aB paralogs. *Genome*. 2011;54(9):738–51.
32. Savada RP, Bonham-Smith PC. Charge versus sequence for nuclear/nucleolar localization of plant ribosomal proteins. *Plant Mol Biol*. 2013;81(4–5):477–93.
33. Lam YW, Lamond AI, Mann M, Andersen JS. Analysis of nucleolar protein dynamics reveals the nuclear degradation of ribosomal proteins. *Curr Biol*. 2007;17(9):749–60.
34. Kruger T, Zentgraf H, Scheer U. Intranucleolar sites of ribosome biogenesis defined by the localization of early binding ribosomal proteins. *J Cell Biol*. 2007;177(4):573–8.
35. Browning KS, Bailey-Serres J. Mechanism of cytoplasmic mRNA translation. *Arabidopsis Book*. 2015;13:e0176.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://www.biomedcentral.com/submissions)

